

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Sorption of Heavy Metals on Clay Minerals and Oxides: A Review

Ifeoma Mary Ugwu and Onyedikachi Anthony Igbokwe

Abstract

Sorption of heavy metals plays a vital role in controlling environmental pollution. Here, we reviewed the sorption of heavy metals such as Ni, Co, Cu, Zn, V, Pb, Hg, In, As, Cd, Cr, Ga, Cs, Mn, V, Eu, Mo, Th, TI and Cr on metal oxides and clay minerals. The mechanism of association between these ions and the host minerals, and the factors controlling their sorption are discussed in detail. Both chemical and empirical methods of describing sorption mechanism are discussed. The sorption processes depend on the pH, metal concentration, ionic strength, temperature, time, adsorbent dosage, type of ion, surface area, type of adsorbent modification and nature of adsorbent. The review confirmed that both metal oxides and clay have capability of sequestering heavy metals, however, combination of both metal oxides and modified clay have enhanced capability of removing heavy metals from aqueous solution. These inorganic adsorbents have the regeneration and recycling potentials and can be used to remediate and sequester economic metals for commercial purposes, however, this needs future investigation.

Keywords: sorption, adsorption, heavy metals, remediation, oxides, clay minerals

1. Introduction

Heavy metal pollution is an environmental problem that has harmful effect on both aquatic and terrestrial environment. These metals are released to the environment through activities such as mining, electroplating and manufacturing of paper and pesticides in form of mine tailings or effluents [1]. They have ability to complex with minerals to form inorganic ligands with variable solubility and acid–base potentials, thus, making their remediation from contaminated land/soil a major concern [2, 3].

Pollution arising from heavy metals poses serious health problems to both human beings and animals [1]. *For example*, low concentration of cadmium in human being can lead to ‘Itai-Itai’ disease, lungs cancer, and kidney and liver problems [4, 5]. Ingestion of water contaminated with high level of Cr can cause ulcer, skin inflammation and pulmonary congestion [6, 7]. Health problems associated with ingestion or inhaling of nickel dust are diarrhoea, renal oedema, nausea, chest pain, pulmonary fibrosis and gastrointestinal ache. Excess of lead in the environment can lead to mental retardation, kidney and nervous disease, cancer [8, 9]. The Cu^{2+} is the most toxic form of copper [10], and health problems

associated with copper are anaemia, headache, kidney damage and death [11, 12]. Large amount of zinc in human body can result to anaemia, fever, vomiting and skin inflammation [13]. Mercury ingestion can lead to damaging of reproductive and respiratory system, kidney and brain [14]. Also, health problem associated with arsenic are gastrointestinal problems, cancer, diabetes, and liver tumours [15]. Large concentration of Mn causes retardation of growth, infertility, fever, eye blindness and muscles [15]. Consumption of water with high concentration of cobalt can lead to heart problems, asthma, damage of thyroid, pneumonia, hair and weight loss, nausea and vomiting, [16]. Depending on the route and length of exposure to vanadium, it can lead to immunological, genotoxic, developmental, reproductive and systemic problems and even death [17]. Exposure to indium via respiration can lead to lung disease, kidney and lungs damage [18, 19]. Long term exposure of gallium may damage the kidney, liver, lungs and bone marrow [20]. Health effects of molybdenum are headache, weight loss, joint or muscle ache, diarrhoea, kidney and liver problems [21]. Exposure to radioactive caesium can lead to cells damage, cancer, diarrhoea, vomiting, nausea and mental abnormality in babies [22]. Exposure to thorium workers can cause lung and liver disease, cancer, liver diseases and death at large exposure [23].

Clay minerals and metal oxides are formed from weathering of primary minerals [24]. These minerals have high surface area and can absorb heavy metals from aqueous and natural environments [25]. Clay minerals, bentonite, sepiolite and palygorskite have been used extensively to remove heavy metals from wastewater and agricultural soil, and the mechanisms of remediation are sorption, precipitation and liming [26]. Heavy metal polluted water is mostly remediated using ion-exchange, adsorption and mechanism [27]. Even though, clay minerals have adsorption affinity for heavy metals, e.g. Cu and Zn, their roles are not significant when compared to iron oxides (e.g. goethite), manganese oxides (e.g. birnessite) and organic materials on a unit mass basis [28]. Comparison of the adsorption potentials of deep sea minerals such as clay minerals, Mn-oxides, Fe-oxides, calcite and apatite reveal that Mn and Fe oxides and oxyhydroxides are crucial phases for scavenging heavy metals [29]. Lead readily sorbs on all phases but has greatest affinity for carbonate fluorapatite, however, Caesium (Cs) has affinity for illite [29]. In sea water, hydrated cations such as Mn^{2+} , Co^{2+} , Ni^{2+} , Zn^{2+} , Cu^{2+} , Ba^{2+} , MnCl^+ and PbCl^+ have strong affinity for Mn-oxide while HVO_4^{2-} , MoO_4^{2-} , PbCO_3 and HAsO_4^{2-} preferentially adsorb onto Fe oxyhydroxide [29].

Sorption techniques have the ability to remove these contaminants via adsorption onto sorbents. However, the type of sorbent to be used for remediation depends on the sorption capacity, sorption pH and distribution coefficient of the contaminant [30]. The review on the sorption of Ni(II) adsorbents indicates that bisorbents are the most effective for removal of the ion from aqueous solution [31]. In addition, Pb and Cd be successfully immobilised with wide range of low cost materials such as metal oxides, animal manure, apatite, lime, biosolids, and biochar [32]. However, the review on the adsorption capabilities of low-cost sorbents such as agricultural waste, household wastes, industrial by-products and sludge indicates that even though these material has great adsorption potentials their particle size, surface area, contact time, initial concentration of ion, adoption dose are not stated by the researchers [33]. This poses great limitation for the use of these adsorbents for remediation. Further review of the sorption of nickel, copper, lead, cadmium, caesium, chromium, europium and thorium on both natural and synthetic materials reveal that more studies on multi-component sorption of these metals are required [34]. Therefore, synthesizing effective and low cost adsorbent that can be recycled for removal of contaminants from aqueous environment will be of high interest

[34]. Similarly, synthesising and discovering new novel methods of using modified clay for environmental remediation is required metals [25].



To this end, this paper presents the review the sorption of heavy metals such as Ni, Co, Cu, Zn, V, Pb, Hg, In, As, Cd, Cr, Ga, Cs, Mn, Eu, Mo, Th, Tl and Cr on both natural and synthetic metal oxides and clay minerals for soil and water remediation. The main aim of this review is to summarise the adsorption mechanism and recent remediation method of single and multicomponent system of these toxic metals using clay minerals and metal oxides. In addition, the advantages of reusability these adsorbents after remediation are discussed.

2. Adsorption mechanism and models


2.1 Adsorption mechanism



The interactions between metallic ions and mineral surfaces in any environment allow the sorption of ions to the solid surface, thereby decreasing the concentration of ions in the aqueous phases below the solubility limits of the solid phase [35]. Determination of partitioning of heavy metals among the solid phase and accurately predicting their mobility in natural environment requires adequate knowledge of the chemical process such as ion exchange, adsorption, surface precipitation, coprecipitation and mineralisation, that govern sorption mechanism [36]. For example, cobalt sorbs to goethite through surface complexation, surface precipitate and structural incorporation depending on the concentration and sorption duration [37].

Surfaces of substances (oxides, organic and inorganic) have surface functional groups or hydroxyl groups of solute ligands [38]. These functional groups are dominant in natural environments as oxides of Al, Fe and Si and in aqueous solution. The surface charge of metallic oxy-hydroxides, phosphate and carbonates is formed through ionisation of surface groups [39]. The pH of surface oxide and oxy-hydroxides changes due to adsorption of protons or ions from solution [40]. Thus, the surface charge set up potential differences between the sorbent and sorbate, thereby influencing the approach of ions towards the surfaces [40]. Consequently, surface hydroxyl groups are responsible for complexation on metal oxides, oxyhydroxides, hydroxides and aluminosilicates. The type of surface functional group influences the sequence of the adsorption reaction, variation in adsorption solution chemistry, electrical properties of the interface and the adsorption ability [41]. The reaction of metals and oxygen atoms at the surface results in the formation of hydroxyl groups (OH^-), which are taken as part of the surface instead of solution. The surface hydroxyl is amphoteric (i.e. they can either accept a proton or donate a proton) as shown below:



Adsorption of metal to the surface takes place via substitution of surface protons (H^+) to form inner sphere complexes:








The surface oxygen, $\equiv\text{SO}^-$ may attract ions from bulk solution to form outer sphere complexes [42]:



Bidentate ligands (Lewis bases, e.g. Cl^- , OH^- , H_2O , NH_3^+) are sorbed to the surface by the replacement of the surface hydroxyl group ($\equiv\text{SOH}$) as follows:



Adsorbed ligands can also take up another metal while metals already adsorbed can attach themselves to another ligand [43].



2.2 Adsorption and Kinetic models

An adsorption isotherm is a relationship between the amount adsorbed and the concentration of the adsorbate at a constant temperature [39]. Both empirical and chemical models are used in describing the adsorption reaction. The former model uses mathematical expressions to interpret adsorption (e.g. Freundlich isotherm, Langmuir isotherm and distribution coefficient) whereas the latter uses a molecular approach to calculate equilibrium constants (e.g. constant capacitance model, triple layer models) [44]. Langmuir isotherm is based on the assumption that only one monolayer is formed during reaction, presence of equivalent sites, immobility of adsorbate and absence of adsorbate-adsorbent interaction [39]. The difference between Langmuir isotherm and Freundlich is that Langmuir isotherm considers maximum sorption whereas Freundlich considers infinity site [39]. However, adsorption isotherms are of limited application when modelling because of the complex and variability of natural environment with heavy metal and existence of sorbents with pH dependent surface charges [39]. Therefore, application of surface complexation models that can integrate both the experimental data and the electrostatic interaction between the ion and the oxide surface is required for modelling the mobility sorption of heavy metal. Computer models such as MINTEQA2, FITEQL, EQLFOR, HYDRAQL and SOILCHEM can calculate both equilibrium speciation and constants [44–46].

Adsorption kinetic model is used to investigate sorption mechanism as well as the rate controlling steps in a batch reaction [34]. Kinetic studies is carried out in a way to vary the initial concentration of adsorbate, temperature, time, sorbent dosage, particle size, pH, type of sorbent and sorbate [34]. It is a continuous measurement of experimental data until equilibrium is attained. The result is tested using kinetic equations to determine the best fit, which will give a better understanding of the sorption mechanism. The adsorption isotherms

Isotherm equations	Kinetic equations
$q_e = \frac{RT}{b} \ln (aC_e)$ Temkin	$\frac{dq}{dt} = k(q_e - q)^2$ Pseudo-second-order model
$\theta = \frac{K_{ad}[M]}{1+K_{ad}[M]}$ Langmuir	$\frac{dq}{dt} = k(q_e - q)$ Lagergren model
$\theta = K_{ad}[M]^n$ Freundlich	$\frac{dq}{dt} = \alpha \exp(-\beta q)$ Elovich model
$q_e = q_{\max} \exp\left(-\left(\frac{RT \ln\left(\frac{C_e}{C_s}\right)}{\beta E_0}\right)^2\right)$ Dubinin-Radushkevich	$\frac{dq}{dt} = k_1 C - K_2 C_s$ Bhattacharya and Venkobachar model
$q_e = q_{\max} \frac{BC_e}{q_{\max}(C_e - C_s) + (1+B-1)(C_e/C_s)}$ Brunauer, Emmett and Teller (BET)	$\frac{dq}{dt} = k_{sorp} C(q_e - q) - k_{des} q$ Adam-Bohart-Thomas
$\log \frac{\theta}{C_0} = \log K_{FH} + n_{FH} \log(1 - \theta)$ Halsey	
$q_e = \frac{K_{RP} C_e}{1 + a_{RP} C_e^{\beta}}$ Redlich-Paterson	
$q_e = q_{\max} \frac{(K_s C_e)^{\gamma}}{(1 + (K_s C_e)^{\gamma})}$ Sip	
$q_e = q_{\max} \frac{b_T C_e}{(1 + (b_T C_e)^{1/n_T})^{n_T}}$ Toth	
θ = fraction of surface coverage; $[M]$ = concentration of ion; C_e = concentration at equilibrium; C_s = solubility of adsorbate at a particular temperature; q_{\max} = saturated monolayer sorption capacity; q_e = the amount of solute adsorbed at equilibrium; q = the amount of solute adsorbed at any given time 't'; C = the concentrations of sorbate in solution at any given time 't'; C_s = the concentrations of sorbate in sorbent at any time 't'.	

Table 1.
Adsorption isotherm and kinetic model equations.

and kinetics commonly used are listed in **Table 1**. In addition, most of the adsorption of ions on clays and oxides are best described by Langmuir, Dubinin-Radushkevich and Freundlich isotherm models and Pseudo-second-order kinetic model [47–53].

Better understanding of sorption mechanism is done using technologies and spectroscopy studies such as Attenuated Total Reflection Infrared [37, 54], Fourier Transform Infrared (FTIR) spectroscopy, extend X-ray absorption fine structure spectroscopy (EXAFS) [55–57], powder X-ray diffraction (XRD), X-ray photoelectron spectroscopy (XPS) and Electron paramagnetic resonance (EPR) [58]. Spectroscopic studies show that copper, nickel, zinc and cobalt bind to Mn(IV) vacancy sites in birnessite to form inner-sphere surface complexes via interlayer triple-corner-sharing or enter the vacancy site to become part of the structure [56]. ATR-FTIR is very useful in distinguishing the outer-sphere and inner-sphere complexes on minerals. Yoon et al., determined of the types and structures of the adsorption complexes formed by oxalate at boehmite (γ -AlOOH)/water and corundum (α -Al₂O₃)/water interfaces using in situ ATR-FTIR spectroscopy and quantum chemical [54]. Their findings suggest that the adsorption mechanism involves loss of protons from aqueous species during ligand-exchange reaction and is useful clue on the transport mechanism and removal of toxic elements in natural water.

3. Investigations of the sorption properties of clay and metal oxides

Clay minerals, e.g. montmorillonite, illite, and kaolinite, are known to have ability of adsorbing heavy metals in their inter layers through ion exchange reaction [25]. They generally have high sorption and expansion properties and are widely

used for removal of contaminants from aqueous solution [25]. Their specific roles for adsorption of heavy metals from aqueous solutions and natural environments are discussed below. These adsorbents and the metals they are capable of removing from aqueous solution are summarised in **Table 2**.

Metals	Adsorbents	Temperature (°C)	Concentration	Optimum recovery pH	Time for maximum adsorption	References
Zn(II) Cu(II) Mn(II) Cd(II) Pb(II) Ni(II)	Montmorillonite	25–26.85	500 mg/L	6–8	180 min	[47, 62, 63]
U(VI) Th (IV)	Jordanian Bentonite	25	100mg/L	3	18 h	[49]
Cu(II) Ag(I)	Verde-Iodo bentonite	Cu: 60 Ag: 20	1.2429 mmol/L	<4.5	2 days	[75]
Th (IV)	Activated bentonite					[76]
In(II)	Chitosan-coated bentonite	25	70 mg/L	4.0	300 min	[77]
Cs(II)	Mixture of iron pillared Layered montmorillonite (80%) and goethite (20%)	15	13.3 mg/L	7.3.	30 min	[50]
Cu(II) Cs(II)	Aluminium-pillared-layered montmorillonites	—	100 mg/L	For Cu 4.0–6.0 and For Cs 3.0–8.0	1.5 h	[51]
Cr (VI))	Magnetite nanoparticles	25	50 mg/L	3.0	3 h	[53]
Mn(II) Ni(II) Cu(II)	Hexagonal birnessite	25	143 mg/L	7–8	24 h– 22 days	[55] [100]
Pb(II) Cu(II) Zn(II) Cd(II)	Hexagonal birnessite Hydrous Mn-oxide (HMO)-coated clay	25	2801 mg/L	4.5 to ~5.5	24 h– 2 weeks	[101] [102]
Hg(II) Cr(III) Pb(II) Cu(II) Zn(II) Ba(II) Ni(II) Mn(II) Cd(II) Ba(II) Ag(I)	Ca-montmorillonite	—	1 N	4.8	16 h	[64]

Metals	Adsorbents	Temperature (°C)	Concentration	Optimum recovery pH	Time for maximum adsorption	References
Eu(III)	Na-montmorillonite	25	1×10^{-3} mol/L	~ 7	8 days	[65]
Cd(II)	Heated ball clay Combination of montmorillonite and humic acid	30	50 mg/L	8 8.5	18 h 24 h	[68] [111] [112]
Pb(II) Cd(II) Ni(II) Cu(II)	Kaolinite	30	160 mg/L	7	30 min	[69]
Cd(II) Co(II) Cu(II) Pb(II) Ni(II) Cr(VI)	Acid-activated form kaolinite	30	250 mg/L	Co and Ni: pH 7.8 Cu and Pb: Ph 5.8 Cd: Ph 10, Cr: pH 7	40–240 min	[70] [71]
Cs(II)	Clay soil with predominantly illite as clay mineral	25	4 mg/L	8	16 h	[72] [73]
Tl(I)	Illite	25	8.39×10^{-6} mol/L	7.18	6 days	[74]
Pb(II) Cu(II) Ni(II)	Synthetic magnesium-aluminium layered double hydroxides modified palygorskite	18–40	500 mg/L	5.0	4.0, 8.0, and 20.0 min, for Pb(II), Cu(II) and Ni(II), respectively	[84]
Pb(II)	Palygorskite-iron oxide nanocomposite	25			12 h	[85]
As(III) Cd(II) Cr(III) Cu(II) Hg(II) Ni(II) Pb(II)	Fe-oxide modified smectites (montmorillonite and saponite)	25	100 µg/L	pH 5	2 h	[88]
Ga (III) V(V) Co(II) Ni(II)	Goethite	20–25	10–58 mg/L	pH 7	24 h	[38] [93] [94] [98]
Cu(II) Zn(II) Cd(II) Pb(II)	Amine-magnetic Fe-oxide	30	1.0 mmol/L	6	60 min	[103]
Ni(II)	Mixture of goethite, humic acid and kaolinite	30	100 mg/L	5.5	2 h	[104]

Table 2.
Summary of metals, potential adsorbents and experimental parameters.

3.1 Montmorillonite clay

The sorption ability of montmorillonite and its remediation potential for nickel and copper from polluted water was investigated by [47]. In their study, sorption rate follows pseudo-second-order rate model and the adsorption model fits to Langmuir model. Also, the point of zero charge (PZC) for the aqueous solution containing both metal at different ionic strengths 0.01 and 0.001 M KCl, is $\text{pH } 3.4 \pm 0.2$ [47]. In contrast, the sorption of cadmium, chromium and lead (II) from wastewater on clay follow 1st order kinetics and can be modelled with both Freundlich and Langmuir model [59]. This study reveals that clay has the capability of removing these metals from aqueous solution at low temperature. In another study, the adsorption of Pb on clay from Tunisia reached maximum at pH 7 and declined above pH 7 due to formation of Pb precipitates, and the mechanism of adsorption via ion exchange [60]. Similarly, sorption of Cu(II) and Ni(II) onto white montmorillonite from obtained Kahramanmaras show increased adsorption with increasing amount of the initial amount of adsorbent, pH and temperature. The white montmorillonite shows more affinity for Cu(II) with 100% adsorption compared to Ni(II) [61]. In another study, sorption of Ni(II) on montmorillonite is influence by the presence of copper in a bi-metal aqueous solution with distortion of the shape of the adsorption isotherm whereas the presence of nickel has negligible effect on the sorption of copper [58]. EPR and XAS analyses of adsorption data reveal that both copper form inner-sphere complexes at pH 8 in both single-metal and binary metal system. However, nickel in single metal system forms inner-sphere complex at pH 8 but in binary-solute systems, it forms Ni phyllosilicate co-precipitate/ $\alpha\text{-Ni(OH)}_2(\text{s})$ precipitate at pH 8.0 [58]. In another study, the sorption of Ni(II) and Mn(II) on montmorillonite from Ugwuoba, Nigeria, reveals 90% removal of these ions from solution containing 500 mg/L at pH 6.0 with maximum adsorption capacity of 166.67 mg/g for nickel and 142.86 mg/g for manganese [62]. Acid treated montmorillonite has enhance adsorption ability for heavy metals (Zn, Cu, Mn, Cd, Pb, Ni), depending on the initial metal concentration [63].

Similarly, the adsorption of mercury, chromium, lead, copper, zinc, barium, nickel, manganese, cadmium, silver on Ca-montmorillonite from single metal solutions followed second-order at greater rate compared to montmorillonite [64]. Copper, lead and chromium forms surface precipitation with montmorillonite whereas mercury, zinc, barium, nickel manganese, cadmium and silver sorb onto montmorillonite via adsorption and ion exchange to form both inner and outer sphere complex at the silanol and aluminol site [64]. However, sorption of bi-metal solution of these metals is better described with Langmuir isotherm model [64]. In another study, Europium (III) sorbs on Na-montmorillonite to form an outer-sphere complex at the exchange site as well as inner-sphere surface complexes at the 'aluminol' and 'silanol' edge sites [65]. In another study, heavy metals such as Hg(II), Cr(III), Pb(II), Cu(II), Zn(II), Ba(II), Ni(II), Mn(II), Cd(II), Ba(II) and Ag (I) in a single metal system adsorbs more on Ca-montmorillonite compared to Na-montmorillonite [66]. The adsorption selective sequence is $\text{Hg(II)} > \text{Zn(II)} > \text{Ba(II)} > \text{Cd(II)} > \text{Ni(II)} > \text{Mn(II)}$. However, Na-Montmorillonite is more effective for removal of Pb(II), Cu(II), Co(II), Cd(II) and Zn(II) from aqueous solution than Ca-Montmorillonite [66]. In general, montmorillonite clay and its modified form has stronger adsorption capacity for arsenic, cadmium, chromium, cobalt, lead, iron, manganese, nickel and zinc than kaolinite and its modified form [67].

3.2 Kaolinite

Adsorption of Cd, Cu, Ni, Zn, Pb and Cr with Ball clay (kaolinite with high plasticity) at 30°C indicates that it sorbs more ion compared to Ca-montmorillonite,

illite, kaolinite and Kaolin, and has the higher affinity for Cd relative to other metals [68]. The sorption was at the maximum at pH 6 and 100% adsorption of 50 mg/L Cd. The sorption rate follows pseudo-first-order kinetics whereas the Langmuir model is used to describe the isotherm model [68]. Similarly, Pb(II), Cd(II), Ni(II) and Cu(II) sorption on kaolinitic clay from Longyan, China, show maximum sorption in 30 min and reduction of Pb concentration from 160.00 to 8.00 mg/L [69]. Another study found that adsorption of Cd(II), Co(II), Cu(II), Pb(II) and Ni(II) onto Acid treated kaolinite have great potential for removal these metals from aqueous solution compared to untreated kaolinite [70]. However, the use of natural and modified kaolinite clays for the removal of Cr(VI) for contaminated water is pH dependent and the adsorption increases from pH 1–2 at equilibration time of 240 min [71].

3.3 Illite

The adsorption of Caesium on clay made up of predominately of illite in the presence of small amount of organic matter and Fe-Oxide indicate that presence of organic matter play significant role for Cs uptake. The adsorption is described by Freundlich [48]. Maximum sorption of Caesium on to clay soil occurs at pH 8 and ambient temperature [72]. Comparative study of Cs sorption on illite, montmorillonite and kaolinite show that Cs is most reactive and has strong affinity for illite compared to other clay [73]. The sorption data was successfully modelled with 1-pK Diffuse layer model [73]. In another study, thallium is found to associate with illite in natural environment and sorption studies reveal that Tl(I) sorbs more onto illite than smectite [74]. The affinity for Tl follows this sequence: $\text{MnO}_2 > \text{illite} > \text{smectite} = \text{ferrihydrite} > \text{Al}_2\text{O}_3 = \text{goethite} > \text{SiO}_2$. However, in presence of Rb, Cs, Tl adsorption to illite is less [74].

3.4 Bentonite

Removal of silver and copper in a binary solution containing the two ions with Verde-lodo bentonite reveals that the adsorbent has high adsorption affinity for copper than silver [75]. Cu adsorption on Verde-lodo bentonite is more at an elevated temperature and attained adsorption capacity of 0.110 mmol/g at 60°C, whereas silver attained 0.090 mmol/g at 20°C [75]. In another study, adsorption of indium on chitosan-coated bentonite is best described by Langmuir isotherm whereas the adsorption kinetics fits the pseudo second order [76]. Thorium (IV) sorption onto activated bentonite depends on temperature, pH, ionic strength, and type of anion and the adsorption kinetics can be described by pseudo-second-order model [77]. The activated bentonite is effective for removal of thorium (IV) and associates with it via surface complexation [77]. In the another study, bentonite was effectively used to remove thorium (IV) at pH 3, equilibration time of 18 h and 25°C and adsorption was fitted to Freundlich, Langmuir and Dubinin-Radushkevich isotherm models. Desorption experiment shows that thorium (IV) is best recovered with 1.0 M HNO_3 [49]. However, the use of bentonite for sorption of heavy metals has some negative effect. This is because increase in concentration of heavy metals such as zinc, lead and copper results to decrease in liquid limit, swelling potential, swelling pressure and free swelling of bentonite but increases its hydraulic conductivity due to sorption of heavy metal in the double layer structure [78].

3.5 Natural clay

Competitive adsorption and desorption cadmium, chromium, copper, lead and zinc on natural clay show difference in the optimum maximum pH of adsorption

and metal affinity for single and multi-element adsorption [79]. The adsorption sequence for single metal sorption is $\text{Cr} > \text{Pb} > \text{Cu} > \text{Cd} > \text{Zn}$ whereas $\text{Cr} > \text{Cu} > \text{Pb} > \text{Cd} > \text{Zn}$ is the obtained sequence in the multicomponent system. The study also notes that the initial pH determines the solubility and removal of heavy metal in competitive multi element scenario [79]. In both the single and multielement adsorption, about 90–100% metal uptake except Zn was achieved at pH 2–12 [79]. In another study, sorption of Cd, Cu, Pb, and Zn with concentration of 10 mmol/L on clay minerals, Fe rich clay minerals, clay-Fe oxide, Fe-oxyhydroxides, and calcite at both alkaline and acidic condition show that cadmium and zinc are adsorbed in acidic media by clay minerals whereas copper and lead sorb more at alkaline media [80]. Sorption of Pb(II), Cd(II), Cu(II) and Zn(II) on clay obtained from Aleg formation, Tunisia that has surface area of $71 \text{ m}^2/\text{g}$ indicate adsorption sequence of $\text{Pb(II)} > \text{Cu(II)} > \text{Zn(II)} > \text{Cd(II)}$ and capacity of 131.58 mg/g in single metal systems and less than 50.10 mg/g for multi-metal [81]. Another study found that Cd(II) adsorption is effective for clay samples heated at 200°C , however, above 200°C adsorption decreases as a result of loss of SiOH or Al–OH binding sites [68]. Similarly, clay from china shows to be effective for removal of Cd at 30°C [82]. In addition, investigation for the temperature and pH effect on Cr (VI) adsorption reveal that sorption rate increases with increasing temperature and decreasing pH for concentration range of $0.743\text{--}1.422 \text{ mg/g}$ [83].

3.6 Palygorskite

Synthetic magnesium-aluminium layered double hydroxides modified palygorskite shows high adsorption capacity for Pb(II), Cu(II) and Ni(II) in the aqueous solution with initial concentration of 100 mg/L at pH 5.0 [84]. The reaction fit into Langmuir isotherm and the pseudo-second-order kinetic model [84]. In another study, the use of synthetic palygorskite-Fe oxide nanocomposite removal of Pb(II) from aqueous solution with initial $20\text{--}500 \text{ mg/L}$ Pb indicates that the adsorbent has maximum Pb(II) adsorption capacity of 26.6 mg/g at 5 g/L adsorbent 150 min^{-1} stirring speed, and 25°C [85]. The adsorption data best fitted to the Langmuir isotherm model and pseudo-second-order kinetic model [85].

3.7 Modified clay

Montmorillonite clay modified with tetramethylammonium cations show enhances adsorption capacity for copper compared with unmodified montmorillonite, and removed up to 99.4% Cu(II) from aqueous solution with adsorption capacity of 925.93 mg/g [86]. Another study found that modification of clay minerals with hexadecylammonium bromide (HDTMA) is effective for attenuation of As(III) and As(V) in the presence of Cd(II), Cu(II) and Mn(II) and anions such as glycine, iminodiacetic acid, ethylenediaminetetraacetic acid disodium salt, oxalic acid, phosphate and sulfate [87]. In contrast, Fe-oxide modified smectites (montmorillonite and saponite) have the best adsorption capacity for removal of As (III), Cd(II), Cr(III), Cu(II), Hg(II), Ni(II), Pb(II) and Zn(II) from aqueous solution with at concentration of $10\text{--}100 \text{ ppb}$ [88]. Mercury is sequestered in Al pillared Clay from Aleg formation, Tunisia at pH 3.2 and 240 min equilibration time [89]. The adsorption fit to Langmuir model with adsorption capacities 49.75 mg/g whereas the adsorption kinetics fits to pseudo-second-order kinetic. The adsorption reaction is exothermic [89]. Another study found that Indium sorbs readily on montmorillonite Metosol modified with di(2-ethylhexyl) phosphoric acid with kinetics that follow pseudo-first- and pseudo-second-order models [89]. Sorption of copper and caesium on Al-pillared montmorillonites follow a pseudo-first-order equation and pH of sorption $4.0\text{--}6.0$ for copper and $3.0\text{--}8.0$ for caesium [51].

Caesium sorption is best described with Langmuir model while copper fits the Freundlich isotherm model [51]. Sorption of Cs onto Fe-pillared montmorillonite, goethite and their mixture show that optimum sorption pH range is 5–9 and 93% removal of Cs occurred with the mixture of 20% goethite with Fe-pillared montmorillonite. The sorption process is exothermic and follows pseudo-first-order kinetics and can be described with Dubinin-Radushkevich, Freundlich and Langmuir models [50]. Also, combination of Al-pillared montmorillonite and Fe₃O₄ an MMT/Fe₃O₄ removes Cs from contaminated soil and water and can be recycled efficiently, and can be applied for the remediation of Cs contaminated land. The adsorption follow pseudo-second-order kinetics and Freundlich isotherm model [52]. In addition, maximum adsorption of Cr(IV) from aqueous solution was achieved by using clay heated at 200 to 400°C [91]. The heating process increased the exposure of the adsorptive site and the adsorptive capabilities [91].

3.8 Metal oxides

The Fe³⁺ oxides and Mn-oxides such as goethite, magnetite and birnessite are usually used in sorption studies of heavy metals and anions because of its simple synthesis in the laboratory [40, 92]. These metal oxides act as hosts for toxic trace elements with valences of +2, +3 and +5 [92]. The potential use of these minerals and their mixtures for remediation of contaminated land are reviewed below.

3.8.1 Goethite

Gallium adsorbs on to goethite to form Ga hexa-coordinated >FeOGa (OH)(2) (0) surface complexes at both acid and alkaline pH and exhibit the same isotope fractionation [93]. In another study, vanadium (V) adsorbs onto goethite to form bidentate corner-sharing complex from initial solution of 2.5 and 25 mg/L at pH 1.5 to 12 [94]. Mononuclear V(V) complexes are present at 25 ppm whereas both mononuclear and polynuclear V(V) complexes exist at 25 mg//solution species form [94]. In another study, at concentration < 10 mg/L and pH 7, cobalt sorbs on goethite at pH 7 to form bidentate edge and corner complexes but form surface precipitates, bidentate edge and corner complexes from 10 to 58 mg/L and pH 7.5 [37]. With ageing, Co sorption on goethite is irreversible depending on the initial concentration [37]. In addition, sorption of Pb, Hg, Cd, Zn, Cu, Ni, Co, Mn, Cr and Al on goethite from solution 10⁻⁶ M was achieved in 2 h and 20°C, but different pH [94]. The pH of 50% sorbed metals (pH 50) ranges from 2.81 for Hg to 6.45 for Mn [95]. The adsorption of Cu(II) with initial concentration of 25 mg/L onto goethite, haematite and lepidocrocite from pH 2–7 reveal that at pH of total metal sorption is 6 for goethite, 6.2 for haematite and 6.8 for lepidocrocite [56]. In same study, EXAFS spectra show that Cu(II) adsorbs these Fe-(hydr)oxides to form two or three bidentate edge sharing, corner sharing and tridentate complexes [56]. Adsorption of Ni, Zn, and Ca onto the goethite reveal that goethite has stronger affinity for Ni and Zn compared to Ca, however, in no observable competition occur in Ni–Ca and Zn–Ca bi-metal systems [96]. The adsorption is well described with single-site Langmuir model [96]. In another study, indium with concentration of 6–29 µg/L was immobilised from acid mine drainage by raising it pH to 8 and subsequent sorption onto iron oxide [97]. In addition, up to 100 mg/L Ni sorbs onto goethite at pH 8, 25°C and equilibration time of 21 days [98].

3.8.2 Birnessite

Mn(II) forms edge-sharing complex with birnessite whereas Ni(II) forms triple corner-sharing complexes at pH 6.5–7.5, however, the addition of Mn(II) in

Ni(II)-birnessite suspension at pH 6.5 results to formation of edge-sharing Ni(II) complexes due to site competition [99]. However, at pH 7.5, the presence of Mn(II) results to transformation of birnessite into feitknechtite that encourages sorption and incorporation of Ni(II) from solution [99]. This suggests that alteration of birnessite can influence the solubility of nickel in anaerobic environment [98]. In another study, hexagonal birnessite (δ -MnO₂) was used to sorb Cu(II) containing 143, 77 and 32 mg/L at pH 1–9 [100]. At pH 3, 100% adsorption results to about 5, 2.5 and 1% of copper in the adsorbate [100]. However, EXFAS characterisation of adsorbates at pH 4 reveals that Cu forms by inner-sphere complexation whereas at pH 8, it associates with birnessite via structural incorporation [100]. Sorption of Zn onto synthetic δ -MnO₂ from an initial solution containing 2000 mg/L of Zn occurs at pH \sim 1 and at maximum pH \sim 5 [101]. However, desorption reaction indicates the Zn sorption is reversible and EXFAS show that it form inner-sphere surface complexes at high pH [101]. Similarly, removal of Pb from solution containing 810, 1782, 2801 mg/L occur at pH \sim 5.5 pH and equilibration time of 2 weeks, thus indication high sorption capacity of birnessite for Pb [101]. Also, desorption experiment show that the sorption is reversible at pH \sim 1 [101]. In another study, sorption of Pb(II), Cu(II), Zn(II), Cd(II) onto hexagonal birnessite was carried out at pH 4.5 and characterised with powder X-ray diffraction (XRD) and X-ray photoelectron spectroscopy (XPS). The result of the study indicates that sorption capacity of birnessite for these metals follow this sequence: $\text{Pb}^{2+} \gg \text{Cu}^{2+} > \text{Zn}^{2+} > \text{Cd}^{2+}$ but Pb sorption is up to 3.9 times more than other metals [102].

3.8.3 Magnetite

Synthetic magnetite nano particles was used to remove Hexavalent chromium (Cr(VI)) from synthetic wastewater containing 0 to 50 mg/L Cr(VI) at pH 3.0 and 30°C. Up to 72% of Cr was removed via adsorption and the isotherm fitted to Freundlich model [53]. In addition, rapid sorption of Cu(II), Zn(II), Cd(II), Pb(II) and Ni(II) on synthetic Amine-magnetic Fe-oxide is influenced by pH, ionic strength and the complexation of amino groups [101]. Maximum adsorption occur at pH 6, ionic strength of 1.0 mmol/L NaCl and duration of 60 min and are best described with Pseudo-second-order kinetic model, Langmuir model [101]. The adsorption affinity of the adsorbent follow this sequence $\text{Pb(II)} > \text{Cu(II)} > \text{Zn(II)} > \text{Cd(II)} > \text{Ni(II)}$ [103].

3.9 Mixture of oxide, clay and/or other materials

Mixture of goethite, humic acid and kaolinite are good adsorbents for lead, cadmium, zinc, nickel and copper and the adsorption is better described with Langmuir and Freundlich isotherm model [104]. However, in a five metal ion system (Quinary), the adsorption of lead, cadmium and nickel are affected negatively by the presence of zinc and copper whereas the presence of lead, cadmium and nickel has synergistic effect on the sorption of zinc and copper [104]. Another study found that low-cost adsorbent Algeria clay that is composed of predominantly montmorillonite and kaolinite has the capacity of adsorbing Cu(II) at pH of 6.5 and 20°C with maximum adsorption capacity of 12.22 mg/g [105]. However, treated Algeria clay under similar condition as the untreated clay has adsorption capacity of 15.40 mg/g [105]. The process of adsorption was spontaneous and exothermic [105]. Sorption of copper (II) ion on palygorskite and sepiolite is enhanced at elevated temperature [106]. The adsorption reactions are endothermically driven but sepiolite sorbs copper spontaneously and has more copper retention capacity compared to palygorskite [106]. In another study, combination of montmorillonite

and ZnO show high capacity for adsorption of Pb and Cu from aqueous solution at wide range of pH [107]. The kinetic of the adsorption reaction follow pseudo-second-order whereas the adsorption isotherm is described by Langmuir isotherm [107]. Arsenic adsorption on goethite, amorphous Fe-hydroxide, and Ti(IV)-Fe(III)-Al(III)-pillared bentonite, clay pillared with titanium (IV), iron (III), and aluminium (III) reveal that amorphous Fe-hydroxide has highest adsorption capacity for arsenate and arsenite [108]. Arsenic is stable at pH 7 and is mobilised at pH 4 and 10 but mostly at acidic condition. Mn-oxides, amorphous iron oxides and clay minerals sequester up to 61% of arsenic [109].

Combination of montmorillonite and humic acid in the ratio of 100:3 is efficient for removal of cadmium at pH 8.5 and contact time of 24 h with adsorption capacity of 18.96 mg/g for cadmium [110]. Another study reveal that the adsorption and desorption of cadmium and copper from montmorillonite, allophane, kaolinite, halloysite reveal that sericite has the highest ability for Cd sorption, however, montmorillonite showed greatest retention for Cd [111]. In addition, all clay types has sorption ability for copper with pH of 50% metal sorbed lower than pH of 50% metal sorbed for cadmium sorption [111]. In another study, the removal of Cu and Zn from aqueous solution by Al-montmorillonite, goethite, kaolinite and their mixtures at room temperature and pH 4 reveal that adsorption is via inner and outer sphere complexation [112]. However, mixing of different mineral for sorption of Cu and Zn retards their removal and decreases the exchange of proton and acid/base potential of the reactive sites [112]. The sorption of Zn onto hydrous Mn-oxide (HMO)-coated clay reveals that the affinity the HMO-coated montmorillonite was greater than that of uncoated montmorillonite, and possess linear isotherm at pH 5–6 [113]. X-ray absorption spectroscopy (XAS) reveal reduction of first shell distance at surface loading of 10^{-3} mol and pH 5–7 due to higher electrostatic attraction [113]. In another study, sorption of copper, zinc and lead on soil composed of clay minerals (smectites and vermiculites), carbonates and Fe-oxide show that Copper and lead has higher sorption capacity and retention compared to Zn [114]. Clay minerals adsorbs more metals than other phases, however, for lead, similar capacity was obtained for Fe-oxides [114]. The presence of carbonate in alkaline condition increases the amount of metal uptake, and the mixture of clay minerals and Fe-oxide enhanced adsorption of the metals [114]. Competitive sorption and desorption of Cd, Cr, Cu, Ni, Pb and Zn by iron oxide, Mn oxides, kaolinite, vermiculite and mica from initial solution of 100 mg/L show that kaolinite and mica has strong affinity and retention capability for Cd; vermiculite, Cu and Zn; iron oxide and Mn-oxide, Pb [115]. Kaolinite has low retention capability for Cu whereas vermiculite and Mn oxide has greatest retention capability of all the metal [115].

In another study, the effect of increase surface area of clay minerals (kaolinite, montmorillonite and illite) through coating with Fe-oxide, organic matter and Al-oxides for adsorption of heavy metals indicate that coating clay increases the surface area of clay minerals with expectation of Aluminium oxide coated montmorillonite and organic matter coated 2:1 phyllosilicates [116]. Another study found that, both amorphous hydrous manganese oxide (HMO) and HMO-coated montmorillonite sorbs Ni and Pb to form inner-sphere complexes with Ni coordinating to vacant site of Mn-oxide structure and Pb forming bidentate corner-sharing complexes [117]. In addition, another study reveal that montmorillonite clay coated with amorphous (hydrous manganese oxide (HMO), birnessite and pyrolusite) has the same surface properties as the coated oxide, however, the surface area of the coated Montmorillonite increases whereas the while the pore size distribution decreased. The HMO- and birnessite-coated clay still retained their pH (point of net zero charge (p_{nzc})) of 2.8 and 3.1, respectively, [118].

3.10 Advantages of using metal oxides and clays for remediation

The use of clay minerals and oxides is more effective than using other materials for remediation of heavy metals from the environment. For example, the use of sepioloite is better for remediation of Cd compared to $\text{Ca}(\text{OH})_2$ [119]. In addition, clay minerals and oxides can be recycled and regenerated for additional remediation use. For example, synthetic palygorskite-Fe oxide exhibited the capability of removing Pb and three cycles reusable potential and retains its magnetic properties for the removal of the heavy metals [85]. Recovery of metals from sorbents can increase the life cycle and long term remediation cost [120]. The estimated zinc that will accumulate in composite substrate in Force Crag Mine is will amount to €7600 in 10 years, however, removal of this substrate after its exhaustion will cost more than €0.8 M whereas recycling the substrate through acid washing is estimated to cost €155,000 [120]. Similarly, oxides and clay minerals can be recycled and reused for remediation of the contaminant. In addition, these sorbents can be used to remediate economic heavy/rare earth metals for commercial and industrial purposes, however, this requires further investigation.

4. Conclusion

Oxides and clay minerals have large surface areas and can sorb heavy metals via adsorption and ion exchange, respectively. Sorption of these toxic metals on oxides and clay minerals reduce their concentration and mobility in aqueous solution and natural environment. This review confirms that sorption is highly dependent on pH, equilibration time, initial concentration of adsorbate, type of adsorbent, temperature, type of adsorbent modification and surface area. In addition, most of the adsorption of ions on clays and oxides are best described by Langmuir models and Pseudo-second-order model. Most remediation techniques employ the use of permeable reactive barriers (PBRs) to react the contaminant from groundwater. These barriers can be Fe-oxides, montmorillonite, Mn oxide-coated montmorillonite, Fe-oxide-coated montmorillonite [121]. The review confirmed that both metal oxides and clay have capability of sequestering heavy metals, however, combination of both metal oxides and modified clay have enhanced capability of removing heavy metals from aqueous solution. These inorganic adsorbents has the regeneration and recycling potentials and can be used to remediate and sequester economic metals for commercial purposes, however, this needs future investigation.

IntechOpen

Author details

Ifeoma Mary Ugwu^{1*} and Onyedikachi Anthony Igbokwe^{2,3}

1 Enugu State University of Science and Technology, Enugu State, Nigeria

2 Ruhr University Bochum, Germany

3 Federal University Ndufe-Alike Ikwo, Ebonyi State, Nigeria

*Address all correspondence to: iu1591@my.bristol.ac.uk

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Carolin CF, Kumar PS, Saravanan A, Joshiba GJ, Naushad M. Efficient techniques for the removal of toxic heavy metals from aquatic environment: A review. *Journal of Environmental Chemical Engineering*. 2017;**5**(3): 2782-2799
- [2] Lee J-C, Pandey BD. Bio-processing of solid wastes and secondary resources for metal extraction - A review. *Waste Management*. 2012;**32**(1):3-18
- [3] Peligro FR, Pavlovic I, Rojas R, Barriga C. Removal of heavy metals from simulated wastewater by in situ formation of layered double hydroxides. *Chemical Engineering Journal*. 2016; **306**:1035-1040
- [4] Demim S, Drouiche N, Aouabed A, Benayad T, Dendene-Badache O, Semsari SC. Assessment of the physiological effects and heavy metal removal using a response surface approach by *L. gibba*. *Ecological Engineering*. 2013;**61**:426-435
- [5] Filipi M. Mechanisms of cadmium induced genomic instability. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*. 2012;**733**: 69-77
- [6] Miretzky P, Cirelli AF. Cr(VI) and Cr(III) removal from aqueous solution by raw and modified lignocellulosic materials: a review. *Journal Hazardous Materials*. 2010;**180**:1-19
- [7] Hu J, Chen C, Zhu X, Wang X. Removal of chromium from aqueous solution by using oxidized multiwalled carbon nanotubes. *Journal Hazardous Materials*. 2009;**162**:1542-1550
- [8] Qu X, Alvarez PJJ, Li Q. Applications of nanotechnology in water and wastewater treatment. *Water Recourses*. 2013;**47**:3931-3946
- [9] Cechinel MAP, Ulson De Souza SMAG, Ulson De Souza AA. Study of lead (II) adsorption onto activated carbon originating from cow bone. *Journal of Cleaner Production*. 2013;**65**: 342-349
- [10] Awual MR, Ismael M, Khaleque MA, Yaita T. Ultra-trace copper (II) detection and removal from wastewater using novel meso-adsorbent. *Journal of Industrial and Engineering Chemistry*. 2015;**20**:2332-2340
- [11] Tang WW, Zeng GM, Gong JL, Liang J, Xu P, Zhang C, et al. Impact of humic/fulvic acid on the removal of heavy metals from aqueous solutions using nanomaterials: a review. *Science of the Total Environment*. 2014;**468–469**:1014-1027
- [12] Zhou YT, Branford-White C, Nie HL, Zhu ML. Removal of Cu²⁺ from aqueous solution by chitosan-coated magnetic nanoparticles modified with α -ketoglutaric acid. *Journal of Colloid and Interface Science*. 2009;**330**:29-37
- [13] Petcu C, Purcar V, Radu AL, Ianchis R, Elvira A, Sarbu A, et al. Removal of zinc ions from model wastewater system using bicopolymer membranes with fumed silica. *Journal of Water Process Engineering*. 2015;**8**:1-10
- [14] Kumari S, Chauhan GS. New cellulose-lysine schiff-base-based sensor-adsorbent for mercury ions. *ACS Applied Materials & Interfaces*. 2014;**6**: 5908-5917
- [15] Ahmaruzzaman M. Industrial wastes as low-cost potential adsorbents for the treatment of wastewater laden with heavy metals. *Advances in Colloid and Interface Science*. 2011;**116**:36-59
- [16] lenntech.com. 1998-2018. Available from: <http://www.lenntech.com/period>

ic/elements/co.htm [Accessed: Jun 10, 2018]

[17] Agency for Toxic Substances and Disease Registry. Toxicological Profile for Vanadium. 2012. Available from: <https://www.atsdr.cdc.gov/toxprofiles/tp58.pdf> [Accessed: Jun 10, 2018]

[18] Cummings KJ, Virji MA, Park JY, Stanton ML, Edwards NT, Trapnell BC, et al. Respirable indium exposures, plasma indium, and respiratory health among indium-tin oxide (ITO) workers: Respirable indium, plasma indium, and respiratory health. *American Journal of Industrial Medicine*. 2016;7:59

[19] New Jersey Department of Health and Senior Services. Hazardous Substances Sheet: Indium. 1999. Available from: <http://nj.gov/health/eoh/rtkweb/documents/fs/1025.pdf> [Accessed: Jun 10, 2018]

[20] New Jersey Department of Health and Senior Services. Hazardous Substances Sheet: Gallium. 2001. Available from: <http://nj.gov/health/eoh/rtkweb/documents/fs/0956.pdf> [Accessed: Jun 10, 2018]

[21] New Jersey Department of Health and Senior Services. Hazardous Substances Sheet: Molybdenum. 1999. Available from: <http://nj.gov/health/eoh/rtkweb/documents/fs/1312.pdf> [Accessed: Jun 10, 2018]

[22] Agency for Toxic Substances and Disease Registry. Public Health Statement for Cesium. 2004. Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp157-c1-b.pdf> [Accessed: Jun 10, 2018]

[23] Agency for Toxic Substances and Disease Registry. Public Health Statement for Thorium. 1990. Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp147-c1-b.pdf> [Accessed: Jun 10, 2018]

[24] Deer WA, Howie RA, Zussman J. *An Introduction to the Rock—Forming Minerals*. London: Longman; 1992

[25] Uddin MK. A review on the adsorption of heavy metals by clay minerals, with special focus on the past decade. *Chemical Engineering Journal*. 2017;308:438-462

[26] Xu Y, Liang X, Xu Y, Qin X, Huang Q, Wang L, et al. Remediation of heavy metal-polluted agricultural soils using clay minerals: A review. *Pedosphere*. 2017;27(2):193-204

[27] Fu FL, Wang Q. Removal of heavy metal ions from wastewaters: A review. *Journal of Environmental Management*. 2011;92(3):407-418

[28] Wang XL, Li Y. Measurement of Cu and Zn adsorption onto surficial sediment components: New evidence for less importance of clay minerals. *Journal of Hazardous Materials*. 2011;189(3):719-723

[29] Koschinsky A, Winkler A, Fritsche U. Importance of different types of marine particles for the scavenging of heavy metals in the deep-sea bottom water. *Applied Geochemistry*. 2003;18(5):693-710

[30] Alby D, Charnay C, Heran M, Prelot B, Zajac J. Recent developments in nanostructured inorganic materials for sorption of cesium and strontium: Synthesis and shaping, sorption capacity, mechanisms, and selectivity-A review. *Journal of Hazardous Materials*. 2018;344:511-530

[31] Raval NP, Shah PU, Shah NK. Adsorptive removal of nickel (II) ions from aqueous environment: A review. *Journal of Environmental Management*. 2016;179:1-20

[32] Mahar A, Wang P, Li RH, Zhang ZQ. Immobilization of lead and cadmium in contaminated soil using

amendments: A review. *Pedosphere*. 2015;**25**(4):555-568

[33] De Gisi S, Lofrano G, Grassi M, Notarnicola M. Characteristics and adsorption capacities of low-cost sorbents for wastewater treatment: A review. *Sustainable Materials and Technologies*. 2016;**9**:10-40

[34] Zhao G, Wu X, Tan X, Wang X. Sorption of heavy metal ions from aqueous solutions: A review. *The Open Colloid Science Journal*. Bentham Open; 2011;**4**:19-31. DOI: 1876-5300/11

[35] Katz LE, Hayes KF. Surface complexation modeling .2. Strategy for modeling polymer and precipitation reactions at high surface coverage. *Journal of Colloid and Interface Science*. 1995;**170**(2):491-501

[36] Fukushi K. Modelling sorption processes of trace elements by earth surface materials. *Journal of Geography-Chigaku Zasshi*. 2017;**126**(3):325-341

[37] Ugwu IM, Sherman DM. Irreversibility of sorption of cobalt to goethite (α -FeOOH) and disparities in dissolution of aged synthetic Co-goethite. *Chemical Geology*. 2017;**467**: 168-176

[38] Stumm W. *Chemistry of Solid-Water Interface*. New York: Wiley; 1992

[39] Langmuir D. *Aqueous Environmental Geochemistry*. London: Prentice Hal; 1997

[40] Appelo CAJ, Geochemistry PD. *Groundwater and Pollution*. Rotterdam: Balkema; 2005

[41] Davis JA, Kent DB. Surface complexation modeling in aqueous geochemistry. *Mineral-Water Interface Geochemistry*. 1990;**23**:177-260

[42] Sherman DM. Surface complexation modelling: Mineral-fluid equilibria at the molecular scale. *Reviews in Mineralogy and Geochemistry*. 2009;**70**: 181-205

[43] White WM. *Geochemistry*. UK: Willey-Blackwell; 2013. 668 pp

[44] Goldberg S. *Adsorption Models Incorporated into Chemical Equilibrium Models*. Chemical Equilibrium and Reaction Models. USA; 1995

[45] Herbelin A, Westall J. *A Computer Program for Determination of Chemical Equilibrium Constants From Experimental Data*. 3.2 ed. Corvallis, OR: Department of Chemistry, Oregon State University; 1996

[46] Sherman DM, Peacock CL, Hubbard CG. Surface complexation of U(VI) on goethite (α -FeOOH). *Geochimica Et Cosmochimica Acta*. 2008;**72**(2):298-310

[47] Ijagbemi CO, Baek M-H, Kim D-S. Montmorillonite surface properties and sorption characteristics for heavy metal removal from aqueous solutions. *Journal of Hazardous Materials*. 2009;**166**(1): 538-546

[48] Dumat C, Cheshire MV, Fraser AR, Shand CA, Staunton S. The effect of removal of soil organic matter and iron on the adsorption of radiocaesium. *European Journal of Soil Science*. 1997; **48**(4):675-683

[49] Khalili FI, Salameh NH, Shaybe MM. Sorption of uranium (VI) and thorium (IV) by Jordanian Bentonite. *Journal of Chemistry*. 2013;**13**:13

[50] Ararem A, Bouras O, Arbaoui F. Adsorption of caesium from aqueous solution on binary mixture of iron pillared layered montmorillonite and goethite. *Chemical Engineering Journal*. 2011;**172**(1):230-236

- [51] Karamanis D, Assimakopoulos PA. Efficiency of aluminum-pillared montmorillonite on the removal of cesium and copper from aqueous solutions. *Water Research*. 2007;**41**(9): 1897-1906
- [52] Zheng X, Dou J, Yuan J, Qin W, Hong X, Ding A. Removal of Cs⁺ from water and soil by ammonium-pillared montmorillonite/Fe₃O₄ composite. *Journal of Environmental Sciences*. 2017;**56**:12-24
- [53] Padmavathy KS, Madhu G, Haseena PV. A study on effects of pH, adsorbent dosage, time, initial concentration and adsorption isotherm study for the removal of hexavalent chromium (Cr (VI)) from wastewater by magnetite nanoparticles. In: *International Conference on Emerging Trends in Engineering, Science and Technology (Icetest—2015)*, Procedia Technology; Vol. 24. 2016. pp. 585-594
- [54] Yoon TH, Johnson SB, Musgrave CB, Brown GE. Adsorption of organic matter at mineral/water interfaces: I. ATR-FTIR spectroscopic and quantum chemical study of oxalate adsorbed at boehmite/water and corundum/water interfaces. *Geochimica et Cosmochimica Acta*. 2004;**68**(22):4505-4518
- [55] Peacock CL, Sherman DM. Sorption of Ni by birnessite: Equilibrium controls on Ni in seawater. *Chemical Geology*. 2007;**238**(1-2):94-106
- [56] Peacock CL, Sherman DM. Copper (II) sorption onto goethite, hematite, and lepidocrocite: A surface complexation model based on ab initio molecular geometries and EXAFS spectroscopy (Vol 68, p. 2623, 2004). *Geochimica et Cosmochimica Acta*. 2005;**69**(21):5141-5142
- [57] Kwon KD, Refson K, Sposito G. Understanding the trends in transition metal sorption by vacancy sites in birnessite. *Geochimica et Cosmochimica Acta*. 2013;**101**:222-232
- [58] Yang S, Ren X, Zhao G, Shi W, Montavon G, Grambow B, et al. Competitive sorption and selective sequence of Cu(II) and Ni(II) on montmorillonite: Batch, modeling, EPR and XAS studies. *Geochimica et Cosmochimica Acta*. 2015;**166**:129-145
- [59] Khan TA, Singh VV. Removal of cadmium (II), lead (II), and chromium (VI) ions from aqueous solution using clay. *Toxicological and Environmental Chemistry*. 2010;**92**(8):1435-1446
- [60] Eloussaief M, Benzina M. Efficiency of natural and acid-activated clays in the removal of Pb(II) from aqueous solutions. *Journal of Hazardous Materials*. 2010;**178**(1-3):753-757
- [61] Demirkiran AR, Acemioglu B, Gonen T. Sorption of copper and nickel ions from solution by clay minerals. *Oxidation Communications*. 2016;**39**(1): 817-829
- [62] Akpomie KG, Dawodu FA, Adebawale KO. Mechanism on the sorption of heavy metals from binary-solution by a low cost montmorillonite and its desorption potential. *Alexandria Engineering Journal*. 2015;**54**(3):757-767
- [63] Akpomie KG, Dawodu FA. Acid-modified montmorillonite for sorption of heavy metals from automobile effluent. *Beni-Suef University Journal of Basic and Applied Sciences*. 2016;**5**:1-12
- [64] de Pablo L, Lourdes Chavez M, Abatal M. Adsorption of heavy metals in acid to alkaline environments by montmorillonite and Ca-montmorillonite. *Chemical Engineering Journal*. 2011;**171**(3):1276-1286
- [65] Kowal-Fouchard A, Drot R, Simoni E, Marmier N, Fromage F, Ehrhardt JJ. Structural identification of europium

- (III) adsorption complexes on montmorillonite. *New Journal of Chemistry*. 2004;**28**(7):864-869
- [66] Chen C, Liu H, Chen T, Chen D, Frost RL. An insight into the removal of Pb(II), Cu(II), Co(II), Cd(II), Zn(II), Ag(I), Hg(I), Cr(VI) by Na(I)-montmorillonite and Ca(II)-montmorillonite. *Applied Clay Science*. 2015;**118**:239-247
- [67] Sen Gupta S, Bhattacharyya KG. Adsorption of heavy metals on kaolinite and montmorillonite: A review. *Physical Chemistry Chemical Physics*. 2012;**14**(19):6698-6723
- [68] Rao RAK, Kashifuddin M. Adsorption studies of Cd(II) on ball clay: Comparison with other natural clays. *Arabian Journal of Chemistry*. 2016;**9**:S1233-S1S41
- [69] Jiang MQ, Jin XY, Lu XQ, Chen ZL. Adsorption of Pb(II), Cd(II), Ni(II) and Cu(II) onto natural kaolinite clay. *Desalination*. 2010;**252**(1-3):33-39
- [70] Bhattacharyya KG, Sen Gupta S. Adsorptive accumulation of Cd(II), Co(II), Cu(II), Pb(II) and Ni(II) ions from water onto kaolinite: Influence of acid activation. *Adsorption Science & Technology*. 2009;**27**(1):47-68
- [71] Bhattacharyya KG, Gupta SS. Adsorption of chromium (VI) from water by clays. *Industrial and Engineering Chemistry Research*. 2006;**45**(45):7232-7240
- [72] Giannakopoulou F, Haidouti C, Chronopoulou A, Gasparatos D. Sorption behavior of cesium on various soils under different pH levels. *Journal of Hazardous Materials*. 2007;**149**:553-556
- [73] Cherif MA, Martin-Garin A, Gerard F, Bildstein O. A robust and parsimonious model for caesium sorption on clay minerals and natural clay materials. *Applied Geochemistry*. 2017;**87**:22-37
- [74] Martin LA, Wissocq A, Benedetti MF, Latrille C. Thallium (Tl) sorption onto illite and smectite: Implications for Tl mobility in the environment. *Geochimica et Cosmochimica Acta*. 2018;**230**:1-16
- [75] Freitas ED, Carmo ACR, Almeida Neto AF, Vieira MGA. Binary adsorption of silver and copper on Verde-lodo bentonite: Kinetic and equilibrium study. *Applied Clay Science*. 2017;**137**:69-76
- [76] Calagui MJC, Senoro DB, Kan C-C, Salvacion JWL, Futralan CM, Wan M-W. Adsorption of indium (III) ions from aqueous solution using chitosan-coated bentonite beads. *Journal of Hazardous Materials*. 2014;**277**:120-126
- [77] Yin Z, Pan D, Liu P, Wu H, Li Z, Wu W. Sorption behavior of thorium (IV) onto activated bentonite. *Journal of Radioanalytical and Nuclear Chemistry*. 2018;**316**(1):301-312
- [78] Dutta J, Mishra AK. Influence of the presence of heavy metals on the behaviour of bentonites. *Environmental Earth Sciences*. 2016;**75**(11)
- [79] Lukman S, Essa MH, Nuthu D, Mu'azu N, Bukhari A, Basheer C. Adsorption and desorption of heavy metals onto natural clay material: Influence of initial pH. *Journal of Environmental Science and Technology*. 2013;**6**:1-15
- [80] Sipos P, Kis VK, Baldzs R, Toth A, Kovacs I, Nemeth T. Contribution of individual pure or mixed-phase mineral particles to metal sorption in soils. *Geoderma*. 2018;**324**:1-8
- [81] Sdiri A, Higashi T, Chaabouni R, Jamoussi F. Competitive removal of heavy metals from aqueous solutions by montmorillonitic and calcareous clays.

Water Air and Soil Pollution. 2012;**223** (3):1191-1204

[82] Sharma YC. Thermodynamics of removal of cadmium by adsorption on an indigenous clay. Chemical Engineering Journal. 2008;**145**(1):64-68

[83] Weng C-H, Sharma YC, Chu S-H. Adsorption of Cr(VI) from aqueous solutions by spent activated clay. Journal of Hazardous Materials. 2008; **155**(1-2):65-75

[84] Yang F, Sun S, Chen X, Chang Y, Zha F, Lei Z. Mg-Al layered double hydroxides modified clay adsorbents for efficient removal of Pb^{2+} , Cu^{2+} and Ni^{2+} from water. Applied Clay Science. 2016; **123**:134-140

[85] Rusmin R, Sarkar B, Tsuzuki T, Kawashima N, Naidu R. Removal of lead from aqueous solution using superparamagnetic palygorskite nanocomposite: Material characterization and regeneration studies. Chemosphere. 2017;**186**: 1006-1015

[86] Medhi H, Bhattacharyya KG. Kinetic and mechanistic studies on adsorption of Cu(II) in aqueous medium onto montmorillonite K10 and its modified derivative. New Journal of Chemistry. 2017;**41**(22): 13533-13552

[87] Lee SM, Lalhmunsiam T, Tiwari D. Porous hybrid materials in the remediation of water contaminated with As(III) and As(V). Chemical Engineering Journal. 2015;**270**:496-507

[88] Franco F, Benitez-Guerrero M, Gonzalez-Trivino I, Perez-Recuerda R, Assiego C, Cifuentes-Melchor J, et al. Low-cost aluminum and iron oxides supported on dioctahedral and trioctahedral smectites: A comparative study of the effectiveness on the heavy metal adsorption from water. Applied Clay Science. 2016;**119**:321-332

[89] Eloussaief M, Sdiri A, Benzina M. Modelling the adsorption of mercury onto natural and aluminium pillared clays. Environmental Science and Pollution Research. 2013;**20**:469-479

[90] Timofeev KL, Maltsev GI, Sviridov AV. Sorption kinetics of indium, iron, and zinc ions on modified montmorillonite. Moscow University Chemistry Bulletin. 2017;**72**(3):128-134

[91] Alemayehu DD, Sk S, Tessema DA. Assessment of the adsorption capacities of fired clay soils from Jimma (Ethiopia) for the removal of Cr(VI) from aqueous solution. Universal Journal of Environmental Research and Technology. 2012;**2**:411-420

[92] Cornell RMSU. The Iron Oxides: Structure, Properties, Reactions, Occurrences and Uses. Germany: John Wiley and Sons; 2003

[93] Yuan W, Saldi GD, Chen J, Zuccolini MV, Birck J-L, Liu Y, et al. Gallium isotope fractionation during Ga adsorption on calcite and goethite. Geochimica et Cosmochimica Acta. 2018;**223**:350-363

[94] Peacock C, Sherman D. Vanadium (V) adsorption onto goethite (α -FeOOH) at p. 1.5 to 12: A surface complexation model based on ab initio molecular geometries and EXAFS spectroscopy. Geochimica et Cosmochimica Acta. 2004;**68**:1723-1733

[95] Fischer L, Bruemmer GW, Barrow NJ. Observations and modelling of the reactions of 10 metals with goethite: Adsorption and diffusion processes. European Journal of Soil Science. 2007; **58**(6):1304-1315

[96] Trivedi P, Axe L, Dyer J. Adsorption of metal ions onto goethite: single-adsorbate and competitive systems. Colloids and Surfaces A—

Physicochemical and Engineering Aspects. 2001;**191**(1–2):107-121

[97] White SJO, Hussain FA, Hemond HF, Sacco SA, Shine JP, Runkel RL, et al. The precipitation of indium at elevated pH in a stream influenced by acid mine drainage. *Science of the Total Environment*. 2017;**574**:1484-1491

[98] Ugwu IM. Sorption of Ni and Co on goethite with application to Ni laterite formation. United Kingdom: University of Bristol; 2015

[99] Lefkowitz JP, Elzinga EJ. Structural alteration of hexagonal birnessite by aqueous Mn(II): Impacts on Ni(II) sorption. *Chemical Geology Journal*. 2017;**466**:524-532

[100] Sherman DM, Peacock CL. Surface complexation of Cu on birnessite (δ -MnO₂): Controls on Cu in the deep ocean. *Geochimica et Cosmochimica Acta*. 2010;**74**(23):6721-6730

[101] Bacon CGD. Surface complexation of Pb and Zn onto birnessite (δ -MnO₂); controls on pollution in soils and groundwater. United Kingdom: University of Bristol; 2014

[102] Wang Y, Feng X, Villalobos M, Tan W, Liu F. Sorption behavior of heavy metals on birnessite: Relationship with its Mn average oxidation state and implications for types of sorption sites. *Chemical Geology Journal*. 2012;**292-293**:25-34

[103] Lin S, Liu LL, Yang Y, Lin KF. Study on preferential adsorption of cationic-style heavy metals using amine-functionalized magnetic iron oxide nanoparticles (MIONPs-NH₂) as efficient adsorbents. *Applied Surface Science*. 2017;**407**:29-35

[104] Unuabonah EI, Olu-Owolabi BI, Adebowale KO. Competitive adsorption of metal ions onto goethite-humic acid-

modified kaolinite clay. *International Journal of Environmental Science and Technology*. 2016;**13**(4):1043-1054

[105] Djebbar M, Djafri F. Adsorption of Cu(II) on natural and treated clays. *Water Quality Research Journal of Canada*. 2016;**51**(1):26-32

[106] Khosravi P, Shirvani M, Bakhtiary S, Shariatmadari H. Energetic and entropic features of Cu(II) sorption equilibria on fibrous clay minerals. *Water Air and Soil Pollution*. 2016:227-354

[107] Sani HA, Ahmad MB, Hussein MZ, Ibrahim NA, Musa A, Saleh TA. Nanocomposite of ZnO with montmorillonite for removal of lead and copper ions from aqueous solutions. *Process Safety and Environmental Protection*. 2017;**109**:97-105

[108] Lenoble V, Bouras O, Deluchat V, Serpaud B, Bollinger JC. Arsenic adsorption onto pillared clays and iron oxides. *Journal of Colloid and Interface Science*. 2002;**255**(1):52-58

[109] Shaw D. Mobility of arsenic in saturated, laboratory test sediments under varying pH conditions. *Engineering Geology*. 2006;**85**(1–2): 158-164

[110] Zeng X, Wang Z, Ji Z, Wei S. A new montmorillonite/humic acid complex prepared in alkaline condition to remove cadmium in waste water. *Polish Journal of Environmental Studies*. 2015;**24**(2):817-821

[111] Itami K, Yanai J. Sorption and desorption properties of cadmium and copper on soil clays in relation to charge characteristics. *Soil Science and Plant Nutrition*. 2006;**52**(1):5-12

[112] Egirani DE, Baker AR, Andrews JE. Copper and zinc removal from aqueous solution by mixed mineral systems—II.

The role of solution composition and aging. *Journal of Colloid and Interface Science*. 2005;**291**(2):326-333

[113] Boonfueng T, Axe L, Xu Y, Tyson TA. The impact of Mn oxide coatings on Zn distribution. *Journal of Colloid and Interface Science*. 2006;**298**(2):615-623

[114] Sipos P, Nemeth T, Kis VK, Mohai I. Sorption of copper, zinc and lead on soil mineral phases. *Chemosphere*. 2008;**73**(4):461-469

[115] Vega FA, Covelo EF, Vazquez JJ, Andrade L. Influence of mineral and organic components on copper, lead, and zinc sorption by acid soils. *Journal of Environmental Science and Health Part A—Toxic/Hazardous Substances & Environmental Engineering*. 2007;**42**(14):2167-2173

[116] Zhuang J, Yu GR. Effects of surface coatings on electrochemical properties and contaminant sorption of clay minerals. *Chemosphere*. 2002;**49**(6):619-628

[117] Boonfueng T, Axe L, Xu Y, Tyson TA. Nickel and lead sequestration in manganese oxide-coated montmorillonite. *Journal of Colloid and Interface Science*. 2006;**303**(1):87-98

[118] Boonfueng T, Axe L, Xu Y. Properties and structure of manganese oxide-coated clay. *Journal of Colloid and Interface Science*. 2005;**281**(1):80-92

[119] Zhu Q-H, Huang D-Y, Zhu G-X, Ge T-D, Liu G-S, Zhu H-H, et al. Sepiolite is recommended for the remediation of Cd-contaminated paddy soil. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science*. 2010;**60**(2):110-116

[120] Bailey MT, Gandy CJ, Jarvis AP. Reducing life-cycle costs of passive mine water treatment by recovery of

metals from treatment wastes. In: Drebenstedt C, Paul M, editors. *Proceedings IMWA 2016, Freiberg/ Germany, Mining Meets Water – Conflicts and Solutions*. 2016

[121] Park Y, Shin WS, Choi S-J. Sorptive removal of cobalt, strontium and cesium onto manganese and iron oxide-coated montmorillonite from groundwater. *Journal of Radioanalytical and Nuclear Chemistry*. 2012;**292**(2):837-852